## Selected References on Salt Marsh Ecology and Restoration.

- Alberts, J.J. and Z. Filip 1998. Metal binding in estuarine humic and fulvic acids: FTIR analysis of humic acid-metal complexes. Environ Technol 19 (9): 923-931.
- Allen, H.H., E.J. Clairain, R.J. Diaz, A.W. Ford, L.F. Junt and B.R. Wells 1978. Habitat development field investigations--Bolivar Peninsula Marsh and upland habitat development site, Galveston, Texas: summary report. U.S. Army Corps Eng. Waterways Experiment Station, Environmental Lab., Vicksburg, MS.
- Allen, H.H. and J.W. Webb, Jr. 1982. Influence of breakwaters on artificial salt marsh establishment on dredged material. Proceedings of the ninth annual conference on wetland restoration and creation., Tampa, FL., 1: 18-35.
- Allen, H.H., S.O. Shirley and J.W. Webb, Jr. 1986. Vegetative stabilization of dredged material in moderate to high wave-energy environments for created wetlands. Proceedings of the thirteenth annual conference on wetland restoration and creation., Hillsborough, FL, 19-35.
- Allen, H. 1988. Biotechnical stabilization of dredged material shorelines., 116-128. In: Beneficial Uses of Dredged Material M.C. Landin (Ed.), Baltimore MD: US Army Corps of Engineers.
- Allen, H.H. and J.W. Webb, Jr. 1993. Bioengineering methods to establish salt marsh on dredged material. Coastal Zone '93, New Orleans, LA, 118-132.
- Allen, E.A., P.E. Fell, M.A. Peck, J.A. Gieg, C.R. Guthke and M.D. Newkirk 1994. Gut contents of common mummichogs, Fundulus heteroclitus L., in a restored impounded marsh and in natural reference marshes. Est. 17 (2): 462-471.
- Ambrose, R.F. and D.J. Meffert 1999. Fish-assemblage dynamics in Malibu Lagoon, a small, hydrologically altered estuary in southern California. Wetlands 19 (2): 327-340.
- Amiaud, B., J.B. Bouzille, F. Tournade and A. Bonis 1998. Spatial patterns of soil salinities in old embanked marshlands in western France. Wetlands 18 (3): 482-494.
- Anonymous 1995. <u>The Galveston Bay Plan. The comprehensive conservation and management plan for the Galveston Bay ecosystem.</u> GBNEP-49, 457pp. Galveston Bay Estuary Program, Webster, TX.
- Anonymous 1998. <u>Habitat Conservation Blueprint</u>, 189pp. Galveston Bay Foundation, Webster, TX.
- Ansede, J.H., P.J. Pellechia and D.C. Yoch 1999. Selenium biotransformation by the salt marsh cordgrass Spartina alterniflora: Evidence for dimethylselenoniopropionate formation. Environ Sci Technol 33 (12): 2064-2069.
- Athnos, D.L. 1993. Compensatory mitigation in the Gulf coast states: Can we achieve "No net loss" of wetlands? Inst. for Coastal and Estuarine Research, Univ. W. Florida,
- Baca, B.J. and T.W. Kana 1986. Methodology for restoring impounded coastal wetlands. Proceedings of the thirteenth annual conference on wetlands restoration and creation., Hillsborough, FL, 36-44.
- Bacic, M.K., S.Y. Newell and D.C. Yoch 1998. Release of dimethylsulfide from dimethylsulfoniopropionate by plant-associated salt marsh fungi. Appl Environ Microbiol 64 (4): 1484-1489.

- Bacic, M.K. and D.C. Yoch 1998. In vivo characterization of dimethylsulfoniopropionate lyase in the fungus Fusarium lateritium. Appl Environ Microbiol 64 (1): 106-111.
- Bagwell, C.E., Y.M. Piceno, A. AshburneLucas and C.R. Lovell 1998. Physiological diversity of the rhizosphere diazotroph assemblages of selected salt marsh grasses. Appl Environ Microbiol 64 (11): 4276-4282.
- Baltz, D.M., C. Rakocinski and J.W. Fleeger 1993. Microhabitat use by marsh edge fishes in a Louisiana estuary. Env. Biol. Fish. 36 109-126.
- Banner, A. 1977. Revegetation and maturation of restored shoreline in Indian River, Florida. Proceedings of the fourth annual conference on restoration of coastal vegetation in Florida., Tampa, FL, 13-42.
- Beeman, S. 1983. Techniques for the creation and maintenance of intertidal saltmarsh wetlands for landscaping and shoreline protection. Proceedings of the tenth annual conference on wetlands restoration and creation., Tampa, FL, 33-43.
- Beeman, S. 1992. Reconnection and restoration of impounded tidal wetlands near Vero Beach, Florida. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL, 19-24.
- Belaire, C.E. and D. Templet 1993. Innovative use of dredged material in Texas benefits endangered whooping cranes. Proceedings of the 20th annual conference on wetlands restoration and creation., Tampa, FL, 1-14.
- Benner, C.S., P.L. Knutson, R.A. Brochu and A.K. Hurme 1982. Vegetative erosion control in an oligahaline environment, Currituck Sound, North Carolina. Wetlands 2 105-117.
- Bennett, A., T.S. Bianchi, J.C. Means and K.R. Carman 1999. The effects of polycyclic aromatic hydrocarbon contamination and grazing on the abundance and composition of microphytobenthos in salt marsh sediments (Pass Fourchon, LA) I. A microcosm experiment. J Exp Mar Biol Ecol 242 (1): 1-20.
- Berger, J.J. (Ed.) 1990. Ecological Restoration in the San Francisco Bay Area. Berkeley, CA: Restoring the Earth, 213 pp.
- Bernstein, P.G. and R.L. Zepp Jr. 1990. Evaluation of selected wetland creation projects authorized through the Corps of Engineers Section 404 program. U.S. Fish and Wildlife Service, Permits/Licenses Branch, Annapolis, MD, 80.
- Bias, M.A. and M.L. Morrison 1999. Movements and home range of salt marsh harvest mice. Southwest Naturalist 44 (3): 348-353.
- Blair, C. 1991. Successful tidal wetland mitigation in Norfolk, VA., 463-476. In: Coastal Wetlands. H.S. Bolton (Ed.), New York: American Society of Civil Engineers.
- Bockelmann, A.C. and R. Neuhaus 1999. Competitive exclusion of Elymus athericus from a high-stress habitat in a European salt marsh. J Ecol 87 (3): 503-513.
- Bolton, H.S. (Ed.) 1991. Coastal Wetlands. New York: American Society of Civil Engineers, 515 pp.
- Bontje, M.P. 1988. The application of science and engineering to restore a salt marsh. Proceedings of the fifteenth annual conference on wetlands restoration and creation., Tampa, FL, 16-23.
- Bontje, M.P. 1991. A successful salt marsh restoration in the New Jersey meadow-lands.

Proceedings of the eighteenth annual conference on wetlands restoration and creation., Tampa, FL, 5-16.

Bortolus, A. and O. Iribarne 1999. Effects of the SW Atlantic burrowing crab Chasmagnathus granulata on a Spartina salt marsh. Mar Ecol Progr Ser 178 79-88.

Boschker, H.T.S., J.F.C. deBrouwer and T.E. Cappenberg 1999. The contribution of macrophyte-derived organic matter to microbial biomass in salt-marsh sediments: Stable carbon isotope analysis of microbial biomarkers. Limnol Oceanogr 44 (2): 309-319.

Bouchard, V., V. Creach, J.C. Lefeuvre, G. Bertru and A. Mariotti 1998. Fate of plant detritus in a European salt marsh dominated by Atriplex portulacoides (L.) Aellen. Hydrobiologia 374 75-87.

Boyd, M.J. 1982. Salt marsh faunas: colonization and monitoring. Wetland restortation and enhancement in California., Hayward, CA, 110.

Boyer, K.E. and J.B. Zedler 1999. Nitrogen addition could shift plant community composition in a restored California salt marsh. Restor Ecol 7 (1): 74-85.

Brewer, J.S., T. Rand, J.M. Levine and M.D. Bertness 1998. Biomass allocation, clonal dispersal, and competitive success in three salt marsh plants. Oikos 82 (2): 347-353.

Brinson, M.M. and R.R. Christian 1999. Stability of Juncus roemerianus patches in a salt marsh. Wetlands 19 (1): 65-70.

Broderick, M.E. and C.J. Dawes 1998. Seasonal photosynthetic and respiratory responses of the red alga Bostrychia tenella (Ceramiales, Rhodophyta) from a salt marsh and mangal. Phycologia 37 (2): 92-99.

Broome, S.W., W.W. Woodhouse Jr. and E.D. Seneca 1975. The relationship of mineral nutrients to growth of *Spartina alterniflora* in North Carolina. II. The effects of N, P, and Fe fertilizers. Soil Science Society American Proceedings 39 301-307.

Broome, S.W., E.D. Seneca and W.W. Woodhouse Jr. 1982. Establishing brackish marshes on graded upland sites in North Carolina. Wetlands 2 152-178.

Broome, S.W., E.D. Seneca and W.W. Woodhouse Jr. 1983. The effects of source, rate and placement of nitrogen and phosphorous fertilizers on growth of *Spartina alterniflora* transplants in North Carolina. Estuaries 6 212-226.

Broome, S.W., E.D. Seneca and W.W. Woodhouse Jr. 1986. Long-term growth and development of transplants of the salt-marsh grass *Spartina alterniflora*. Estuaries 9 63-74

Broome, S.W., E.D. Seneca and W.W. Woodhouse Jr. 1988. Tidal salt marsh restoration. Aquatic Botany 32 1-22.

Broome, S.W. 1989. Creation and restoration of tidal wetlands of the southeastern United States., 37-72. In: Wetland Creation and Restoration: The Status of the Science. Part 1, Regional Review. J.A. Kusler and M.E. Kentula (Ed.), Washington, D.C.: Island Press, Inc.

Broome, S.W., S.M. Rogers Jr. and E.D. Seneca 1992. Shoreline erosion control using marsh vegetation and low-cost structures.

Bush, J.K. and O.W. VanAuken 1997. The effects of neighbors and grazing on the growth of Helianthus paradoxus. Southwest Naturalist 42 (4): 416-422.

Cai, W.J. and Y. Wang 1998. The chemistry, fluxes, and sources of carbon dioxide in the

estuarine waters of the Satilla and Altamaha Rivers, Georgia. Limnol Oceanogr 43 (4): 657-668.

Callaway, J.C. and M.N. Josselyn 1992. The introduction and spread of smooth cordgrass (*Spartina alterniflora*) in south San Francisco Bay. Estuaries 15 218-226.

Callaway, J.C., R.D. Delaune and W.H. Patrick 1998. Heavy metal chronologies in selected coastal wetlands from Northern Europe. Mar Pollut Bull 36 (1): 82-96.

Callaway, R.M. and S.C. Pennings 1998. Impact of a parasitic plant on the zonation of two salt marsh perennials. Oecologia 114 (1): 100-105.

Cammen, L.M. 1975. Accumulation rate and turnover time of organic carbon in a salt marsh sediment. Limnology and Oceanography 20 1012-1015.

Cammen, L.M. 1976. Abundance and production of macroinvertebrates from natural and artificially established salt marshes in North Carolina. amn 96 487-493.

Cammen, L.M. 1976. Macroinvertebrate colonization of *Spartina* marshes artificially established on dredge spoil. ecms 4 357-372.

Cammen, L.M. 1979. The macro-infauna of a North Carolina salt marsh. American Midland Naturalist 102 244-253.

Cantero, J.J., R. Leon, J.M. Cisneros and A. Cantero 1998. Habitat structure and vegetation relationships in central Argentina salt marsh landscapes. Plant Ecol 137 (1): 79-100.

Carlson, D.B., P.D. OBryan and J.R. Rey 1999. Florida's salt-marsh management issues: 1991-98. J Amer Mosquito Contr Assn 15 (2): 186-193.

Cartaxana, P. and D. Lloyd 1999. N-2, N2O and O-2 profiles in a Tagus estuary salt marsh. Estuar Coast Shelf Sci 48 (6): 751-756.

Castellanos, E.M., C. Heredia, M.E. Figueroa and A.J. Davy 1998. Tiller dynamics of Spartina maritima in successional and non-successional Mediterranean salt marsh. Plant Ecol 137 (2): 213-225.

Chabreck, R.H. 1989. Creation, restoration, and enhancement of marshes of the northcentral Gulf coast., 127-144. In: Wetland creation and restoration: The status of the science. Volume I: Regional overviews. EPA/600/3-89/038. J.A. Kusler and M.E. Kentula (Ed.), Corvallis, OR: U.S. Environmental Protection Agency.

Chamberlain, R.H. and R.A. Barnhart 1993. Early use by fish of a mitigation salt marsh, Humbolt Bay, California. Est. 16 (4): 769-783.

Chapman, H.F., P.E.R. Dale and B.H. Kay 1998. A method for assessing the effects of runneling on salt marsh grapsid crab populations. J Amer Mosquito Contr Assn 14 (1): 61-68.

Chelius, M.K. and J.E. Lepo 1999. Restriction fragment length polymorphism analysis of PCR-amplified nifH sequences from wetland plant rhizosphere communities. Environ Technol 20 (8): 883-889.

Clark, P.A. 1992. Tidal marsh restoration using volunteer fishermen. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL, 36-41.

Cobb, R.A. 1987. Mitigation evaluation study for the south Texas coast, 1975-1986. Corpus Christi State University and U.S. Fish and Wildlife Service,

- Cochran, J.K., M. Frignani, M. Salamanca, L.G. Bellucci and S. Guerzoni 1998. Lead-210 as a tracer of atmospheric input of heavy metals in the northern Venice Lagoon. Mar Chem 62 (1-2): 15-29.
- Coen, L.D., M.W. Luckenbach and D.L. Breitburg 1998. The role of oyster reefs as essential fish habitat: A review of current knowledge and some new perspectives, 438-454. In: Fish Habitat: Essential Fish L.R. Benaka (Ed.), 5410 Grosvenor Lane, Ste 110, Bethesda, MD 20814-2199, USA: Amer Fisheries Soc.
- Courtney, F.X., S.A. Peck and M.O. Hall 1991. Post-harvest recovery of a donor *Spartina alterniflora* marsh. Proceedings of the eighteenth annual conference on wetlands restoration and creation., Tampa, FL, 23-31.
- Covin, J.D. and J.B. Zedler 1988. Nitrogen effects on *Spartina foliosa* and *Salicornia virginica* in the salt marsh at Tijuana Estuary, California. Wetlands 8 51-66.
- Craft, C.B., S.W. Broome and E.D. Seneca 1988. Nitrogen, phosphorus and organic carbon pools in natural and transplanted marsh soils. Estuaries 11 272-280.
- Craft, C.B., E.D. Seneca and S.W. Broome 1991. Porewater Chemistry of Natural and Created Marsh Soils. Journal of Exploratory Marine Biology and Ecology 152 187-200.
- Crewz, D.W. and R.R. Lewis III 1991. An evaluation of historical attempts to establish emergent vegetation in marine wetlands in Florida. Gainesville, FL: Florida Sea Grant, Univ. of Florida,, 113.
- Cross, R.E. and A.E. Stiven 1999. Size-dependent interactions in salt marsh fish (Fundulus heteroclitus Linnaeus) and shrimp (Palaemonetes pugio Holthuis). J Exp Mar Biol Ecol 242 (2): 179-199.
- Cruse, V.L. 1993. An evaluation of the effectiveness of wetland mitigation in Broward County, Florida permitted by the Florida Department of Environmental Regulation. Proceedings of the 20th annual conference on wetlands restoration and creation., Tampa, FL, 45-65.
- Currin, C.A. and H.W. Pearl 1998. Epiphytic nitrogen fixation associated with standing dead shoots of smooth cordgrass, Spartina alterniflora. Estuaries 21 (1): 108-117.
- Dale, P.E.R., S.A. Ritchie, H. Chapman and M.D. Brown 1999. Eggshell sampling: Quantitative or qualitative data? J Amer Mosquito Contr Assn 15 (1): 74-76.
- Day, J.W., F. Scarton, A. Rismondo and D. Are 1998. Rapid deterioration of a salt marsh in Venice Lagoon, Italy. J Coastal Res 14 (2): 583-590.
- Devai, I., R.D. DeLaune, C.B. Henry, C.W. Lindau and P.O. Roberts 1998. Environmental significance of atmospheric emission resulting from in situ burning of oiled salt marsh. Mar Environ Res 45 (2): 157-167.
- Drent, R.H. and R. vanderWal 1999. Cyclic grazing in vertebrates and the manipulation of the food resource, 271-299. In: Herbivores: Between Plants an H. Olff, V.K. Brown and R.H. Drent (Ed.), Osney Mead, Oxford Ox2 0EI, England: Blackwell Science Publ.
- Earhart, H.G. and E.W. Garbisch Jr. 1983. Habitat development utilizing dredged material at Barren Island, Dorchester County, Maryland. Wetlands 3 108-119.
- Earhart, H.G. and E.W. Garbisch Jr. 1986. Beneficial uses of dredged materials at Barren Island, Dorchester County, Maryland. Proceedings of the thirteenth annual conference on wetlands restoration and creation., Hillsborough, FL, 75-85.

Edgcomb, V.P., J.H. McDonald, R. Devereux and D.W. Smith 1999. Estimation of bacterial cell numbers in humic acid-rich salt marsh sediments with probes directed to 16S ribosomal DNA. Appl Environ Microbiol 65 (4): 1516-1523.

Edwards, R.E. 1994. Fishery value of a Sarasota Bay, Florida habitat enhancement project. Proceedings of the 21st annual conference on wetlands restoration and creation., Tampa, FL, 33-48.

Eleuterius, L.N. and J.I. Gill 1981. Long-term observations on seagrass beds and salt marsh established from transplants. Proceedings of the eighth annual conference on wetlands restoration and creation., Tampa, FL., 74-86.

Emmerson, R.H.C., S.B. OReillyWiese, C.L. Macleod and J.N. Lester 1997. A multivariate assessment of metal distribution in inter-tidal sediments of the Blackwater Estuary, UK. Mar Pollut Bull 34 (11): 960-968.

Espey Huston & Associates Inc 1988. Monitoring of transplanted *Spartina alterniflora* on an unconfined dredged material disposal site, Chocolate Bay, Texas. Espey, Huston and Associates, Inc..

Espey Huston & Associates Inc 1990. Monitoring of transplanted *Spartina alterniflora* on an unconfined dredged material disposal site, Pelican Spit, Galveston Bay, Texas. Espey, Huston & Associates, Inc.,

Esselink, P., K.S. Dijkema, S. Reents and G. Hageman 1998. Vertical accretion and profile changes in abandoned man-made tidal marshes in the Dollard estuary, the Netherlands. J Coastal Res 14 (2): 570-582.

Faber, P., A. Shepherd and P. Williams 1988. Monitoring a tidal restoration site in San Francisco Bay- the Muzzi Marsh. Urban wetlands: proceedings of the national wetland symposium., Oakland, CA, 331-335.

Faber, P.M. 1991. The Muzzi marsh, Corte Madera, California: long-term observations of a restored marsh in San Francisco Bay., 424-438. In: Coastal Wetlands. H.S. Bolton (Ed.), New York: American Society of Civil Engineers.

Florida Department of Environmental Regulation. 1991. Report on the effectiveness of permitted mitigation. Florida Department of Environmental Regulation.,

Ford, M.A., D.R. Cahoon and J.C. Lynch 1999. Restoring marsh elevation in a rapidly subsiding salt marsh by thin-layer deposition of dredged material. Ecol Eng 12 (3-4): 189-205.

Fowler, B.K., G.R. Hardaway, G.R. Thomas, C.L. Hill, J.E. Frye and N.A. Ibison 1985. Vegetative growth patterns in planted marshes of the vegetative erosion control project. Proceedings of the twelth annual conference on wetlands restoration and creation., Tampa, FL, 110-120.

Frenkel, R.E. and L.M. Kunze 1984. Introduction and spread of three *Spartina* species in the Pacific Northwest. Annual Meeting of the Association of American Geographers, Washington, DC,

Gallagher, J.L. 1980. Salt marsh soil development. Rehabilitation and creation of selected coastal habitats: proceedings of a workshop, FWS/OBS-80/27: 28-34.

Gallagher, J.L., G.F. Somers, D.M. Grant and D.M. Seliskar 1988. Persistent differences in two forms of *Spartina alterniflora*: A common garden experiment. Ecology 69(4) 1005-1008.

Garbisch, E.W., Jr., P.B. Woller and R.J. McCallum 1975. Salt marsh establishment and development. Fort Belvoir, VA.: U.S. Army Corps Eng., Coastal Eng. Resch. Cent., 114.

Gerard, V.A. 1999. Positive interactions between cordgrass, Spartina alterniflora, and the brown alga, Ascophyllum nodosum ecad scorpioides, in a mid-Atlantic coast salt marsh. J Exp Mar Biol Ecol 239 (1): 157-164.

Goodfriend, W.L. 1998. Microbial community patterns of potential substrate utilization: A comparison of salt marsh, sand dune, and seawater-irrigated agronomic systems. Soil Biol Biochem 30 (8-9): 1169-1176.

Gregg, C.S. and J.W. Fleeger 1998. Grass shrimp Palaemonetes pugio predation on sediment- and stem-dwelling meiofauna: field and laboratory experiments. Mar Ecol Progr Ser 175 77-86.

Hacker, S.D. and M.D. Bertness 1999. Experimental evidence for factors maintaining plant species diversity in a New England salt marsh. Ecology 80 (6): 2064-2073.

Halpin, P.M. and K.L.M. Martin 1999. Aerial respiration in the salt marsh fish Fundulus heteroclitus (Fundulidae). Copeia (3): 743-748.

Hardaway, C.S., G.R. Thomas, B.K. Fowler, C.L. Hill, J.E. Frye and N.A. Ibison 1985. Results of the vegetative erosion control project in the Virginia Chesapeake Bay system. Proceedings of the twelth annual conference on wetlands restoration and creation., Tampa, FL, 144-158.

Hartman, R.D., R.N. Ruebsamen, P.M. Jones and J.L. Koellen 1993. The National Marine Fisheries Service habitat conservation efforts in Louisiana, 1980 through 1990. Mar. Fish. Rev. 54 11-20.

Hartman, R.D., T. Osborn, B. Julius, M. Newell, A. Arnold and E. Zobrist 1994. Mitigating an oil spill in Timbalier Bay, Louisiana: NOAA's Damage Assessment and Restoration Program in action. Proceedings of the 21st annual conference on wetlands restoration and creation., Tampa, FL, 61-76.

Hemminga, M.A., J. vanSoelen and Y.E.M. Maas 1998. Biomass production in pioneer Spartina anglica patches: Evidence for the importance of seston particle deposition. Estuar Coast Shelf Sci 47 (6): 797-805.

Hettler Jr., W.F. 1989. Nekton use of regularly-flooded saltmarsh cordgrass habitat in North Carolina, USA. Marine Ecology Progress Series 56 111-118.

Hoback, W.W., T.M. Svatos, S.M. Spomer and L.G. Higley 1999. Trap color and placement affects estimates of insect family-level abundance and diversity in a Nebraska salt marsh. Entomol Exp Appl 91 (3): 393-402.

Hoffman, W.E. and J.A. Rodgers 1980. A cost/benefit analysis of two large coastal plantings in Tampa Bay, Florida. Proceedings of the seventh annual conference on wetlands restoration and creation., Tampa, FL, 265-278.

Hogan, T.J. and J.R. Ingram 1992. Creating wetland mitigation partnerships. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL, 63-73.

Holman, R.E. 1994. A survey of wetland restoration and creation practitioners. Proceedings of the 21st annual conference on wetlands restoration and creation., Tampa, FL, 77-86.

Horne, M.T., N.J. Finley and M.D. Sprenger 1999. Polychlorinated biphenyl- and

- mercury-associated alterations on benthic invertebrate community structure in a contaminated salt marsh in southeast Georgia. Arch Environ Contam Toxicol 37 (3): 317-325.
- Hovel, K.A. and S.G. Morgan 1999. Susceptibility of estuarine crab larvae to ultraviolet radiation. J Exp Mar Biol Ecol 237 (1): 107-125.
- Hsieh, Y.P., C.H. Yang and J.N. Feng 1998. Sulfate reduction and a molybdate-induced soluble nitrogen flush in sediments during incubation. Soil Biol Biochem 30 (13): 1799-1804.
- Hughes, C.E., P. Binning and G.R. Willgoose 1998. Characterisation of the hydrology of an estuarine wetland. J Hydrol 211 (1-4): 34-49.
- Huisman, J. and H. Olff 1998. Competition and facilitation in multispecies plant-herbivore systems of productive environments. Ecol Lett 1 (1): 25-29.
- Huisman, J., J.P. Grover, R. vanderWal and J. vanAndel 1999. Competition for light, plant-species replacement and herbivore abundance along productivity gradients, 239-269. In: Herbivores: Between Plants an H. Olff, V.K. Brown and R.H. Drent (Ed.), Osney Mead, Oxford Ox2 0El, England: Blackwell Science Publ.
- Josselyn, M.J. (Ed.) 1982. Wetland restoration and enhancement in California. La Jolla, CA: California Sea Grant College Program, 110 pp.
- Josselyn, M.N., J.B. Zedler and T. Griswold 1989. Wetland mitigation along the Pacific coast of the United States., 1-36. In: Wetland creation and restoration: The status of the Science. Part 1, Regional review. J.A. Kusler and M.E. Kentula (Ed.), Washington, DC: Island Press.
- Kentula, M.E., R.P. Brooks, S.E. Gwin, C.C. Holland, A.D. Sherman and J.C. Sifneos 1992. An Approach to Improving Decision Making in Wetland Restoration and Creation. Corvallis, OR: U.S. Environmental Protection Agency, Environmental Research Laboratory, 151.
- Khan, H.R., D.A. Price, K.E. Doble, M.J. Greenberg and A.S.M. Saleuddin 1999. Osmoregulation and FMRFamide-related peptides in the salt marsh snail Melampus bidentatus (Say) (Mollusca: Pulmonata). Biol Bull 196 (2): 153-162.
- Kiraly, S.J., F.A. Cross and J.D. Buffington 1991. The federal effort to evaluate coastal wetland mitigation: A report by the National Ocean Pollution Policy board's habitat loss and modification working group. NOAA Technical Memorandum CS/NOPPO 91-2. 10 p. plus Appendices.,
- Kirchner, G. and H. Ehlers 1998. Sediment geochronology in changing coastal environments: Potentials and limitations of the Cs-137 and Pb-210 methods. J Coastal Res 14 (2): 483-492.
- Klap, V.A., P. Louchouarn, J.J. Boon, M.A. Hemminga and J. vanSoelen 1999. Decomposition dynamics of six salt marsh halophytes as determined by cupric oxide oxidation and direct temperature-resolved mass spectrometry. Limnol Oceanogr 44 (6): 1458-1476.
- Knutson, P.L., J.C. Ford, M.R. Inskeep and J. Oyler 1981. National survey of planted salt marshes (vegetative stabilization and wave stress). Wetlands 1 129-157.
- Knutson, P.L., R.A. Brochu, W.N. Seelig and M. Inskeep 1982. Wave damping in *Spartina alterniflora* marshes. Wetlands 2 87-104.

- Kohlmeyer, J., H.O. Baral and B. VolkmannKohlmeyer 1998. Fungi on Juncus roemerianus. 10. A new Orbilia with ingoldian anamorph. Mycologia 90 (2): 303-309.
- Kraus, D.B. and M.L. Kraus 1986. The establishment of a fiddler crab (*Uca minax*) colony on a manmade *Spartina* mitigation marsh, and its effect on invertebrate colonization. National wetland symposium: mitigation of impacts and losses, Berne, NY, 343-348.
- Kruczynski, W.L. 1982. Salt marshes of the northeastern Gulf of Mexico., 71-88. In: Creation and restoration of coastal plant communities R.R. Lewis, III (Ed.), Boca Raton, FL: CRC Press, Inc.
- Kuhn, N.L., I.A. Mendelssohn and D.J. Reed 1999. Altered hydrology effects on Louisiana salt marsh function. Wetlands 19 (3): 617-626.
- Kusler, J.A. and M.E. Kentula (Ed.) 1989. Wetland Creation and Restoration. The Status of the Science. Washington, DC: Island Press, 595.
- Laffaille, P., S. Brosse, E. Feunteun, A. Baisez and J.C. Lefeuvre 1998. Role of fish communities in particulate organic matter fluxes between salt marshes and coastal marine waters in the Mont Saint-Michel Bay. Hydrobiologia 374 121-133.
- Landin, M.C. and J.W. Webb, Jr. 1986. Wetland development and restoration as part of Corps of Engineer programs: Case studies. National Wetland Symposium: Mitigation of Impacts and Losses., New Orleans, LA, 388-391.
- Langis, R., M. Zalejko and J.B. Zedler 1991. Nitrogen assessments in a constructed and a natural salt marsh of San-Diego Bay. Ecological Applications 1 40-51.
- LaPerriere, A.J. and M.M. Farmer 1989. Recent wetland restoration/creation actions in the New York District. Proceedings of the sixteenth annual conference on wetlands restoration and creation., Plant City, FL., pp 97-108.
- LaSalle, M.W., M.C. Landin and J.G. Sims 1991. Evaluation of the flora and fauna of a *Spartina alterniflora* marsh established on dredged material in Winyah Bay, South Carolina. Wetlands 11 191-208.
- LaSalle, M.W. 1996. Assessing the functional level of a constructed intertidal marsh in Mississippi. Technical Report, WRP-RE-15. U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Lee, C.R. 1992. Creation of wetlands with dredged material: Pros and cons. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL. 86-90.
- Levine, J.M., S.D. Hacker, C.D.G. Harley and M.D. Bertness 1998. Nitrogen effects on an interaction chain in a salt marsh community. Oecologia 117 (1-2): 266-272.
- Levine, J.M., J.S. Brewer and M.D. Bertness 1998. Nutrients, competition and plant zonation in a New England salt marsh. J Ecol 86 (2): 285-292.
- Lewis, R.R., III and C.S. Lewis 1977. Tidal marsh creation on dredged material in Tampa Bay, Florida. Proceedings of the fourth annual conference on restoration of coastal vegetation in Florida., Tampa, FL, 45-67.
- Lewis, R.R., III 1989. Wetlands restoration/creation/enhancement terminology: suggestions for standardization., 417-419. In: Wetland creation and restoration: The status of the Science. Part 2. Perspectives. J.A. Kusler and M.E. Kentula (Ed.), Washington, DC: Island Press.
- Lewis, R.R., III 1989. Creation and restoration of coastal plain wetlands in Florida., 73-

- 102. In: Wetland creation and restoration: The status of the Science. Part 1, Regional review. J.A. Kusler and M.E. Kentula (Ed.), Washington, DC: Island Press.
- Lewis, R.R., III 1992. Coastal habitat restoration as a fishery management tool., In: Stemming the tide of coastal fish habitat loss. R.H. Stroud (Ed.), Savannah: National Coalition for Marine Conservation, Inc.
- Liebezeit, G., T.D. Kunnemann and G. Gad 1999. Biotechnological potential of North Sea salt marsh plants a review of traditional knowledge. J Biotechnol 70 (1-3): 77-84.
- Lin, Q., I.A. Mendelssohn, C.B. Henry, P.O. Roberts, M.M. Walsh, E.B. Overton and R.J. Portier 1999. Effects of bioremediation agents on oil degradation in mineral and sandy salt marsh sediments. Environ Technol 20 (8): 825-837.
- Lindall, W.N., Jr. and G.W. Thayer 1982. Quantification of National Marine Fisheries Service habitat conservation efforts in the southeast region of the United States. Mar. Fish. Rev. 44 18-22.
- Lindau, C.W. and L.R. Hossner 1981. Substrate characterization of an experimental marsh and three natural marshes. Soil Science, American Journal 45 1171-1176.
- Lindau, C.W. and L.R. Hossner 1982. Sediment fractionation of Cu, Ni, Zn, Cr, Mn, and Fe in one experimentl and three natural marshes. Journal of Environmental Quality 11 540-545.
- Loidi, J., M. Herrera, I. Biurrun and I. GarciaMijangos 1999. Relationships between syntaxonomy of Thero-Salicornietea and taxonomy of the genera Salicornia and Suaeda in the Iberian Peninsula. Folia Geobot 34 (1): 97-114.
- Lowe, G., R.J. Epting and L.M. Grant 1993. Wetland creation: A viable mitigation option. Proceedings of the 20th annual conference on wetlands restoration and creation., Tampa, FL, 155-163.
- Lyon, J.T., III 1975. A comparison of epiphytes on natural and planted *Spartina alterniflora* marshes. M.S. Thesis, North Carolina State University, 48.
- Mager, A. and G.W. Thayer 1986. NMFS habitat conservation efforts in the Southeast Region of the United States from 1981 through 1985. Marine Fisheries Review 48 1-8.
- Maranon, T. 1998. Soil seed bank and community dynamics in an annual-dominated Mediterranean salt-marsh. J Veg Sci 9 (3): 371-378.
- Markley, S.M., G.R. Milano and E. Calas 1992. Biscayne Bay restoration and enhancement program shoreline and habitat enhancement guide. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL, 111-120.
- Matthews, G.A. and T.J. Minello 1994. Technology and success in restoration, Creation, and enhancement of *Spartina alterniflora* marshes in the United States. Volume 1 -- Executive summary and annotated bibliography. Decision Analysis Series No. 2, 71 pp.
- Matthews, G.A. and T.J. Minello 1994. Technology and success in restoration, Creation, and enhancement of *Spartina alterniflora* marshes in the United States. Volume 2 -- Inventory and human resources directory. Decision Analysis Series No. 2, 352 pp.
- McTigue, T.A. and R.J. Zimmerman 1998. The use of infauna by juvenile Penaeus aztecus Ives and Penaeus setiferus (Linnaeus). Estuaries 21 (1): 160-175.
- Meyer, D.L., M.S. Fonseca, D.R. Colby, W.J. Kenworthy and G.W. Thayer 1993. An examination of created marsh and seagrass utilization by living marine

resources, 1858-1863. In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management. O. Magoon, W.S. Wilson, H. Converse and L.T. Tobin (Ed.), New York.: American Society Of Civil Engineers.

Middelburg, J.J., J. Nieuwenhuize, R.K. Lubberts and O. vandePlassche 1997. Organic carbon isotope systematics of coastal marshes. Estuar Coast Shelf Sci 45 (5): 681-687.

Miller, L., G.T. Auble and K.A. Schneller-McDonald 1991. User's guide to the wetland creation/restoration data base, version 2. Wash. D.C.: U.S. Dept. Inter., Fish and Wildlife Service, Research and Development,

Miller, J.A. 1993. Juvenile chinook (Oncorhynchus tshawytscha) and coho (O. kisutch) salmon in natural and created estuarine habitats: foraging and daily growth. Univ. of Washington.

Miller, D.L., F.E. Smeins and J.W. Webb 1998. Response of a Texas distichlis spicata coastal marsh following lesser snow goose herbivory. Aquat Bot 61 (4): 301-307.

Minello, T.J., R.J. Zimmerman and E.F. Klima 1987. Creation of fishery habitat in estuaries., 106-117. In: Beneficial uses of dredged material; Proceedings of the first Interagency Workshop, 7-9 Oct 1986, Pensacola, Florida. M.C. Landin and H.K. Smith (Ed.), US Army COE, WES: Tech. Rept. D-87-1.

Minello, T.J. and R.J. Zimmerman 1992. Utilization of natural and transplanted Texas salt marshes by fish and decapod crustaceans. Marine Ecology Progress Series 90 273-285.

Minello, T.J. and J.W. Webb, Jr 1993. The development of fishery habitat value in created salt marshes, 1864-1865. In: Coastal Zone '93, Volume 2. Proceedings of the 8th Symposium on Coastal and Ocean Management. O. Magoon, W.S. Wilson, H. Converse and L.T. Tobin (Ed.), New York: American Society Of Civil Engineers.

Minello, T.J., R.J. Zimmerman and R. Medina 1994. The importance of edge for natant macrofauna in a created salt marsh. wetlands 14 (3): 184-198.

Minello, T.J. and J.W. Webb Jr. 1997. Use of natural and created Spartina alterniflora salt marshes by fishery species and other aquatic fauna in Galveston Bay, Texas, USA. Marine Ecology Progress Series 151 165-179.

Morris, J.T. and A. Jensen 1998. The carbon balance of grazed and non-grazed Spartina anglica saltmarshes at Skallingen, Denmark. J Ecol 86 (2): 229-242.

Morrison, J. and P. Williams 1988. Warm Springs marsh restoration. Urban wetlands: proceedings of the national wetland symposium., Oakland, CA, 340-349.

Moy, L.D. and L.A. Levin 1991. Are *Spartina* marshes a replaceable resource? A functional approach to evaluation of marsh creation efforts. Estuaries 14 1-16.

Mulder, C.P.H. and R.W. Ruess 1998. Relationships between size, biomass allocation, reproduction, and survival in Triglochin palustris: implications for the effects of goose herbivory. Can J Bot 76 (12): 2164-2176.

Mulder, C.P.H. and R.W. Ruess 1998. Effects of herbivory on arrowgrass: Interactions between geese, neighboring plants, and abiotic factors. Ecol Monogr 68 (2): 275-293.

Munro, J.W. 1991. Wetland restoration in the mitigation context. Restoration & Management Notes 9 80-86.

Naidoo, G. and Y. Naidoo 1998. Salt tolerance in Sporobolus virginicus: the importance of ion relations and salt secretion. Flora 193 (4): 337-344.

Nailon, R.W. and E.L. Seidensticker 1991. The effects of shoreline erosion in Galveston Bay, Texas, 193-206. In: Coastal wetlands H.S. Bolton (Ed.), New York: Amer. Soc. Civil Engineers.

National Research Council 1992. Restoration of Aquatic Ecosystems: Science, Technology, and Public Policy. Washington D.C.: National Academy Press, 552.

Netto, S.A. and P.C. Lana 1999. The role of above- and below-ground components of Spartina alterniflora (Loisel) and detritus biomass in structuring macrobenthic associations of Paranagua Bay (SE, Brazil). Hydrobiologia 400 167-177.

Newling, C.J., M.C. Landin and S.D. Parris 1983. Long-term monitoring of the Apalachicola Bay wetland habitat development site. Proceedings of the tenth annual conference on wetland restoration and creation., Tampa, FL, 164-186.

Nicolau, B.A. and J.W. Tunnell, Jr. 1996. Estuarine faunal use in a mitigation project, Nueces River delta, Texas: Sixth year report with a review from 1989 through 1995. Corpus Christi: Texas A&M Univ.-Corpus Christi, 105 pp.

Niering, W.A. 1989. Vegetation dynamics in relation to wetland creation., 479-486. In: Wetland creation and restoration: The status of the Science. Part 2. Perspectives. J.A. Kusler and M.E. Kentula (Ed.), Washington, DC: Island Press.

Nomann, B.E. and S.C. Pennings 1998. Fiddler crab-vegetation interactions in hypersaline habitats. J Exp Mar Biol Ecol 225 (1): 53-68.

Nordlie, F.G. and D.C. Haney 1998. Adaptations in salt marsh teleosts to life in waters of varying salinity. Ital J Zoology 65 405-409.

Orson, R.A., R.S. Warren and W.A. Niering 1998. Interpreting sea level rise and rates of vertical marsh accretion in a southern New England tidal salt marsh. Estuar Coast Shelf Sci 47 (4): 419-429.

Pacific• Estuarine Research Laboratory 1990. A manual for assessing restored and natural coastal wetlands with examples from southern California. California Sea Grant Report No. T-CSGCP-021. La Jolla, California: 105.

Padgett, D.E., C.B. Rogerson and C.T. Hackney 1998. Effects of soil drainage on vertical distribution of subsurface tissues in the salt marsh macrophyte Spartina alterniflora Lois. Wetlands 18 (1): 35-41.

Padgett, D.E. and J.L. Brown 1999. Effects of drainage and soil organic content on growth of Spartina alterniflora (Poaceae) in an artificial salt marsh mesocosm. Amer J Bot 86 (5): 697-702.

Page, H.M. 1997. Importance of vascular plant and algal production to macro-invertebrate consumers in a southern California Salt Marsh. Estuar Coast Shelf Sci 45 (6): 823-834.

Paludan, C. and J.T. Morris 1999. Distribution and speciation of phosphorus along a salinity gradient in intertidal marsh sediments. Biogeochemistry 45 (2): 197-221.

Pantoja, S. and C. Lee 1999. Peptide decomposition by extracellular hydrolysis in coastal seawater and salt marsh sediment. Mar Chem 63 (3-4): 273-291.

Parsons, M.L. 1998. Salt marsh sedimentary record of the landfall of Hurricane Andrew on the Louisiana coast: Diatoms and other paleoindicators. J Coastal Res 14 (3): 939-950.

Paterson, A.W. and A.K. Whitfield 1997. A stable carbon isotope study of the food-web in a freshwater-deprived South African estuary, with particular emphasis on the ichthyofauna. Estuar Coast Shelf Sci 45 (6): 705-715.

Patience, N. and V.V. Klemas 1993. Wetland functional health assessment using remote sensing and other techniques: literature search. NOAA, 114.

Pennings, S.C. and C.L. Richards 1998. Effects of wrack burial in salt-stressed habitats: Batis maritima in a southwest Atlantic salt marsh. Ecography 21 (6): 630-638.

Pennings, S.C., T.H. Carefoot, E.L. Siska, M.E. Chase and T.A. Page 1998. Feeding preferences of a generalist salt-marsh crab: Relative importance of multiple plant traits. Ecology 79 (6): 1968-1979.

Person, B.T., C.A. Babcock and R.W. Ruess 1998. Forage variation in brood-rearing areas used by Pacific black brant geese on the Yukon-Kuskokwim delta, Alaska. J Ecol 86 (2): 243-259.

Peterson, G.W. and R.E. Turner 1994. The value of salt marsh edge vs interior as a habitat for fish and decapod crustaceans in a Louisiana tidal marsh. Est. 17 235-262.

Piceno, Y.M., P.A. Noble and C.R. Lovell 1999. Spatial and temporal assessment of diazotroph assemblage composition in vegetated salt marsh sediments using denaturing gradient gel electrophoresis analysis. Microbial Ecol 38 (2): 157-167.

Piehler, M.F., C.A. Currin, R. Cassanova and H.W. Paerl 1998. Development and N-2-fixing activity of the benthic microbial community in transplanted Spartina alterniflora marshes in North Carolina. Restor Ecol 6 (3): 290-296.

Pinkney, A.E., P.C. McGowan, D.R. Murphy, T.P. Lowe, D.W. Sparling and W.H. Meredith 1999. Effects of temephos (Abate (R) 4E) on fiddler crabs (Uca pugnax and Uca minax) on a delaware salt marsh. J Amer Mosquito Contr Assn 15 (3): 321-329.

Pontee, N.I., J.P. Tastet and L. Masse 1998. Morpho-sedimentary evidence of Holocene coastal changes near the mouth of the Gironde and on the Medoc Peninsula, SW France. Oceanol Acta 21 (2): 243-261.

Portnoy, J.W., B.L. Nowicki, C.T. Roman and D.W. Urish 1998. The discharge of nitrate-contaminated groundwater from developed shoreline to marsh-fringed estuary. Water Resour Res 34 (11): 3095-3104.

Portnoy, J.W. 1999. Salt marsh diking and restoration: Biogeochemical implications of altered wetland hydrology. Environ Manage 24 (1): 111-120.

Posey, M.H., T.D. Alphin and C.M. Powell 1997. Plant and infaunal communities associated with a created marsh. Estuaries 20 (1): 42-47.

Quintana, X.D., R. MorenoAmich and F.A. Comin 1998. Nutrient and plankton dynamics in a Mediterranean salt marsh dominated by incidents of flooding. Part 1: Differential confinement of nutrients. J Plankton Res 20 (11): 2089-2107.

Quintana, X.D., F.A. Comin and R. MorenoAmich 1998. Nutrient and plankton dynamics in a Mediterranean salt marsh dominated by incidents of flooding. Part 2: Response of the zooplankton community to disturbances. J Plankton Res 20 (11): 2109-2127.

Rand, T.A. 1999. Effects of environmental context on the susceptibility of Atriplex patula to attack by herbivorous beetles. Oecologia 121 (1): 39-46.

Reppert, R. 1992. Wetlands mitigation banking concepts. U.S. Army Corps of

Engineers, 33.

Reubsamen, R.N. 1986. National Marine Fisheries Service efforts in Texas to tailor mitigation to specific wetland types. Proceedings of the National Wetland Symposium: Mitigation of Impacts and Losses., New Orleans, LA, 1: 424-425.

Riggs, S. 1992. Distribution of *Spartina alterniflora* in Padilla Bay, Washington, in 1991. Washington Dept. Ecology, Padilla Bay National Estuarine Research Reserve, 63.

Roberts, T.H. 1989. Habitat value of man-made coastal marshes in Florida. Sixteenth annual conference on wetlands restoration and creation., Plant City, FL., 157-179.

Roberts, T.H. 1991. Habitat value of man-made coastal marshes in Florida. Vicksburg, MS: U.S. Army Corps Eng., Waterways Experiment Station, 42.

Rogers, J., J. Harris and I. Valiela 1998. Interaction of nitrogen supply, sea level rise, and elevation on species form and composition of salt marsh plants. Biol Bull 195 (2): 235-237.

Roman, C.T., J.A. Peck, J.R. Allen, J.W. King and P.G. Appleby 1997. Accretion of a New England (USA) salt marsh in response to inlet migration, storms, and sea-level rise. Estuar Coast Shelf Sci 45 (6): 717-727.

Rowcliffe, J.M., A.R. Watkinson and W.J. Sutherland 1998. Aggregative responses of brent geese on salt marsh and their impact on plant community dynamics. Oecologia 114 (3): 417-426.

Rozas, L.P. and T.J. Minello 1998. Nekton use of salt marsh, seagrass, and nonvegetated habitats in a south Texas (USA) estuary. Bull Mar Sci 63 (3): 481-501.

Rozsa, R. and R.A. Orson 1993. Restoration of degraded salt marshes in Connecticut. Proceedings of the 20th annual conference on wetlands restoration and creation., Tampa, FL, 196-205.

Sacco, J. 1989. Infaunal community development of artificially established salt marshes in North Carolina. Ph.D. Thesis, North Carolina State University., 41.

Sacco, J.N., E.D. Seneca and T.R. Wentworth 1994. Infaunal community development of artificially established salt marshes in North Carolina. Est. 17 (2): 489-500.

Sakolsky, G., D.A. Carlson, B.D. Sutton and J.G. Stoffolano 1999. Detection of cryptic species in the Tabanus nigrovittatus (Diptera: Tabanidae) complex in Massachusetts by morphometric and cuticular hydrocarbon analysis. J Med Entomol 36 (5): 610-613.

Sanderson, E.W., M. Zhang, S.L. Ustin and E. Rejmankova 1998. Geostatistical scaling of canopy water content in a California salt marsh. Landscape Ecol 13 (2): 79-92.

Sanger, D.M., A.E. Holland and G.I. Scott 1999. Tidal creek and salt marsh sediments in South Carolina coastal estuaries: I. Distribution of trace metals. Arch Environ Contam Toxicol 37 (4): 445-457.

Sanger, D.M., A.F. Holland and G.I. Scott 1999. Tidal creek and salt marsh sediments in South Carolina coastal estuaries: II. Distribution of organic contaminants. Arch Environ Contam Toxicol 37 (4): 458-471.

Sarda, R., K. Foreman, C.E. Werme and I. Valiela 1998. The impact of epifaunal predation on the structure of macroinfaunal invertebrate communities of tidal saltmarsh creeks. Estuar Coast Shelf Sci 46 (5): 657-669.

- Seliskar, D.M. 1998. Natural and tissue culture-generated variation in the salt marsh grass Sporobolus virginicus: Potential selections for marsh creation and restoration. Hortscience 33 (4): 622-625.
- Seneca, E.D., S.W. Broome and W.W. Woodhouse Jr. 1985. Comparison of *Spartina alterniflora* Loisel. transplants from different locations in a man-initiated marsh in North Carolina. Wetlands 5 181-190.
- Seneca, E.D., S.W. Broome and W.W. Woodhouse 1985. The influence of duration-of-inundation on development of a man-initiated <u>Spartina</u> <u>alterniflora</u> Loisel marsh in North Carolina. J. Exp. Mar. Biol. Ecol. 94 259-268.
- Seneca, E.D. and S.W. Broome 1992. Restoring tidal marshes in North Carolina and France, 53-78. In: Restoring the nation's marine environment G.W. Thayer (Ed.), College Park: Maryland Sea Grant College.
- Sherwood, R.M. and P.S. Petraitis 1998. Mortality differences of two intertidal mussels, Mytilus edulis L. and Geukensia demissa (Dillwyn), in a New Jersey salt marsh. J Exp Mar Biol Ecol 231 (2): 255-265.
- Shirley, M.A. 1992. Recolonization of a restored red mangrove habitat by fish and macroinvertebrates. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL, 159-173.
- Shisler, J.K. 1989. Creation and restoration of the coastal wetlands of the northeastern United States., 145-174. In: Wetland creation and restoration: The status of the Science. Volume I: Regional overviews. EPA/600/3-89/038. J.A. Kusler and M.E. Kentula (Ed.), Corvallis: U.S. Environ. Prot. Agency.
- Sinicrope, T.L., P.G. Hine, R.S. Warren and W.A. Niering 1990. Restoration of an impounded salt marsh in New England. Estuaries 13 25-30.
- Soil Conservation Service 1992. Wetland restoration, enhancement, or creation., 79. In: Engineering Field Handbook Soil Conservation Service (Ed.), Washington DC: U.S. Department of Agriculture.
- Staniforth, R.J., N. Griller and C. Lajzerowicz 1998. Soil seed banks from coastal subarctic ecosystems of Bird Cove, Hudson Bay. Ecoscience 5 (2): 241-249.
- Steinke, T.J. 1986. Hydrologic manipulation and restoring wetland values: Pine Creek, Fairfield, Connecticut. National wetland symposium: mitigation of impacts and losses., New Orleans, LA, 377-383.
- Stout, J.P. 1990. Evaluation of coastal wetlands mitigation, Alabama. Alabama Dept. Environ. Management, 45.
- Streever, W.J. and T.L. Crisman 1992. The effect of inter-wetland and intra-wetland variability on sampling strategies in faunal studies comparing natural and constructed wetlands. Proceedings of the 19th annual conference on wetlands restoration and creation., Tampa, FL, 183-196.
- Streever, W.J., A.J. Genders and M.A. Cole 1998. A closed chamber CO2 flux method for estimating marsh productivity. Aquat Bot 62 (1): 33-44.
- Stumpf, R.P. and J.W. Haines 1998. Variations in tidal level in the Gulf of Mexico and implications for tidal wetlands. Estuar Coast Shelf Sci 46 (2): 165-173.
- Suk, N.S., Q. Guo and N.P. Psuty 1999. Suspended solids flux between salt marsh and adjacent bay: A long-term continuous measurement. Estuar Coast Shelf Sci 49 (1): 61-

81.

Sun, Y. and T. Torgersen 1998. Rapid and precise measurement method for adsorbed Ra-224 on sediments. Mar Chem 61 (3-4): 163-171.

Sundby, B., C. Vale, I. Cacador, F. Catarino, M.J. Madureira and M. Caetano 1998. Metal-rich concretions on the roots of salt marsh plants: Mechanism and rate of formation. Limnol Oceanogr 43 (2): 245-252.

Tanner, G.W. and J.D. Dodd 1985. Effects of phenological stage of *Spartina alterniflora* transplant culms on stand development. Wetlands 4 57-74.

Taylor, M.H. 1999. A suite of adaptations for intertidal spawning. Amer Zool 39 (2): 313-320.

Thayer, G.W. (Ed.) 1992. Restoring the Nation's Marine Environment. College Park, MD: University of Maryland Sea Grant College Program and DOC/NOAA/National Marine Fisheries Service, 716.

Theodose, T.A. and J.B. Roths 1999. Variation in nutrient availability and plant species diversity across forb and graminoid zones of a Northern New England high salt marsh. Plant Ecol 143 (2): 219-228.

Tyler, A.N. 1999. Monitoring anthropogenic radioactivity in salt marsh environments through in situ gamma-ray spectrometry. J Environ Radioact 45 (3): 235-252.

Tyler, A.C. and J.C. Zieman 1999. Patterns of development in the creekbank region of a barrier island Spartina alterniflora marsh. Mar Ecol Progr Ser 180 161-177.

Ugo, P., A. Bertolin and L.M. Moretto 1999. Monitoring sulphur species and metal ions in salt-marsh pore-waters by using an in-situ sampler. Int J Environ Anal Chem 73 (2): 129-143

Underwood, S.G., G.D. Steyer, B. Good and D. Chambers 1991. Bay bottom terracing and vegetative planting: an innovative approach for habitat and water quality enhancement. Proceedings of the eighteenth annual conference on wetland restoration and creation., Hillsborough, FL, 164-173.

Ungar, I.A. 1998. Are biotic factors significant in influencing the distribution of halophytes in saline habitats? Bot Rev 64 (2): 176-199.

US Army Corps of Engineers 1989. Engineering and Design. Environmental Engineering for Coastal Protection. Washington DC: U.S. Army Corps of Engineers, VanAuken, O.W. and J.K. Bush 1998. Spatial relationships of Helianthus paradoxus (Compositae) and associated salt marsh plants. Southwest Naturalist 43 (3): 313-320.

vanderWal, R., P. Kunst and R. Drent 1998. Interactions between hare and brent goose in a salt marsh system: evidence for food competition? Oecologia 117 (1-2): 227-234.

vanWijnen, H.J. and J.P. Bakker 1999. Nitrogen and phosphorus limitation in a coastal barrier salt marsh: the implications for vegetation succession. J Ecol 87 (2): 265-272.

Walsh, M.R., M.E. Walsh and C.M. Collins 1999. Remediation methods for white phosphorus contamination in a coastal salt marsh. Environ Conserv 26 (2): 112-124.

Wang, Y.P., R.S. Zhang and S. Gao 1999. Velocity variations in salt marsh creeks, Jiangsu, China. J Coastal Res 15 (2): 471-477.

Watzin, M.C. and J.G. Gosselink 1992. The fragile fringe: coastal wetlands of the

- continental United States. Louisiana Sea Grant College Program, Louisiana State University, Baton Rouge, LA; U.S. Fish and Wildlife Service, Washington, DC; and National Oceanic and Atmospheric Administration, Rockville, MD.,
- Webb, J.W., Jr. 1977. Establishment of vegetation for shoreline stabilization in Galveston Bay, Texas. Ph.D., Texas A&M Univ., College Station, TX, Webb, J.W., Jr. 1982. Salt marshes of the western Gulf of Mexico., 89-110. In: Creation and restoration of coastal plant communities R.R. Lewis III (Ed.), Boca Raton, FL: CRC Press, Inc.
- Webb, J.W., Jr. and J.D. Dodd 1983. Wave-protected versus unprotected transplantings on a Texas bay shoreline. Journal of Soil & Water Conservation 38 363-366.
- Webb, J.W., Jr. 1983. Soil water salinity variations and their effects on *Spartina alterniflora*. Contributions in Marine Sciience 26 1-13.
- Webb, J.W., Jr., J.D. Dodd, A.T. Weichert and B.H. Koerth 1985. *Spartina alterniflora* response to fertilizer rates, planting dates, and elevation in Galveston Bay, Texas. Estuaries. Second water quality and wetlands management conference., New Orleans, LA, 379-399.
- Webb, J.W., Jr. and C.J. Newling 1985. Comparison of natural and man-made salt marshes in Galveston Bay complex, Texas. Wetlands 4 75-86.
- Webb, J.W., Jr., M.C. Landin and H.H. Allen 1986. Approaches and techniques for wetlands development and restoration of dredged material disposal sites. National wetland symposium: mitigation of impacts and losses., New Orleans, LA, 132-134.
- Webb, J.W., Jr. and J.D. Dodd 1989. *Spartina alterniflora* response to fertilizer, planting dates, and elevation in Galveston Bay, Texas. Wetlands 9 61-72.
- Wiggins, J. and E. Binney 1987. A baseline study of the distribution of *Spartina alterniflora* in Padilla Bay. Washington Dept. Ecology, Padilla Bay National Estuarine Research Reserve, 28.
- Wijte, A.H.B.M. and J.L. Gallagher 1991. Role of live and dead shoots in overwintering Spartina. Bot. Gaz. 152 (4): 509-513.
- Williams, S.L. and J.B. Zedler 1992. Restoring sustainable coastal ecosystems on the Pacific coast: Establishing a research agenda. California Sea Grant College, 19.
- Wilsey, B.J., K.L. McKee and I.A. Mendelssohn 1992. Effects of increased elevation and macro- and micronutrient additions on *Spartina alterniflora* transplant success in saltmarsh dieback areas in Louisiana. Environmental Management 16 505-511.
- Wilson, D.J., R. vanderWal, E.R. Chang, A. Jensen and R.L. Jefferies 1999. Urea hydrolysis and nitrification in arctic salt-marsh soils: Possible constraints on the growth of forage plants. Ecoscience 6 (1): 72-78.
- Wolf, R.B., L.C. Lee and R.R. Sharitz 1986. Wetland creation and restoration in the United States from 1970 to 1985: An annotated bibliography. Wetlands 6 1-88.
- Woodhouse, W.W., Jr. 1979. Building salt marshes along the coasts of the continental United States. Fort Belvoir, VA: U.S. Army Corps Eng., Coastal Eng. Res. Cent., 96.
- Woodhouse, W.W., Jr. and P.L. Knutson 1982. Atlantic coastal marshes., 45-70. In: Creation and restoration of coastal plant communities R.R. Lewis, III (Ed.), Boca Raton, FL: CRC Press, Inc.

Wu, J.L. and D.M. Seliskar 1998. Salinity adaptation of plasma membrane H+-ATPase in the salt marsh plant Spartina patens: ATP hydrolysis and enzyme kinetics. J Exp Bot 49 (323): 1005-1013.

Yang, S.L. 1998. The role of Scirpus marsh in attenuation of hydrodynamics and retention of fine sediment in the Yangtze Estuary. Estuar Coast Shelf Sci 47 (2): 227-233.

Zedler, J.B., M. Josselyn and C. Onuf 1982. Restoration techniques, research, and monitoring: vegetation. Wetland Restoration and Enhancement in California., Hayward, CA., 110.

Zedler, J.B., J. Covin, C. Nordby, P. Williams and J. Boland 1986. Catastrophic events reveal the dynamic nature of salt-marsh vegetation in southern California. Ecology 9 75-80.

Zedler, J.B. and R. Langis 1991. Comparisons of constructed and natural salt marshes of San Diego Bay. Res. Manag. Notes 9 21-25.

Zedler, J.B. 1993. Canopy architecture of natural and planted cordgrass marshes: selecting habitat evaluation criteria. Ecological Applications 3 123-138.

Zeff, M.L. 1999. Salt marsh tidal channel morphometry: Applications for wetland creation and restoration. Restor Ecol 7 (2): 205-211.